

IN THE CLAIMS:

Kindly rewrite Claims 1-51 as follows:

1-20. (Canceled).

21. (Currently Amended) A method of forming an implantable medical device configured to release at least one therapeutic agent therefrom, wherein the therapeutic agent is disposed in a matrix affixed to the body of the implantable medical device, the body includes at least one recess, and wherein the concentration of the at least one therapeutic agent in the matrix varies as a continuous gradient relative to a surface of the body of the implantable medical device, the method comprising:

forming a first homogeneous solution comprising the at least one therapeutic agent mixed with a polymeric binder;

~~applying-introducing~~ the first homogeneous solution into a-the at least one recess ~~in to~~ the body of the implantable medical device;

solidifying the first homogeneous solution, thereby forming a first portion of the matrix;

forming a second homogeneous solution comprising the polymeric binder;

applying the second homogeneous solution to the first portion of the matrix, thereby at least partially liquifying the first portion of the matrix; and

solidifying the second homogeneous solution, thereby forming a second portion of the matrix, wherein the concentration of the at least one therapeutic agent in the matrix is different in the first and second portions of the matrix.

22. (Original) The method of Claim 21, wherein the first and second homogenous solutions include a solvent and the first and second homogenous solutions are solidified by evaporation of the solvent.

23. (Original) The method of Claim 22, wherein the solvent comprises a miscible organic solvent.

24. (Original) The method of Claim 23, wherein the miscible organic solvent is selected from the group consisting of dimethyl sulfoxide, N-methyl pyrrolidone, ethyl lactate, and simple alcohols.

25. (Original) The method of Claim 22, wherein the solvent and the polymeric binder are both non-water soluble.

26. (Original) The method of Claim 25, wherein the non-water soluble solvent is selected from the group consisting of a poly(lactide-co-glycolide) polymer, N-methyl pyrrolidone, ethyl lactate, anisole, chloroform, tetrahydrofuran, xylene, and methylene chloride.

27. (Original) The method of Claim 21, further comprising:
applying successive homogeneous solutions to the matrix; and
solidifying the successive homogeneous solutions, thereby forming additional portions of the matrix, wherein the concentration of the therapeutic agent in the matrix is different in the successive portions of the matrix.

28. (Canceled)

29. (Currently Amended) The method of Claim 21, wherein said at least one recess extends through the body of the implantable medical device~~applying the first homogeneous solution to the implantable medical device body comprises:~~
introducing the homogeneous solution into an opening passing through the ~~implantable medical device body.~~

30. (Canceled)

31. (Original) The method of Claim 21, wherein the polymeric binder of the first homogeneous solution is water soluble.

32. (Original) The method of claim 21, wherein the therapeutic agent is selected from the group consisting of antithrombotic agents , antiproliferative agents, and antirestenotic agents.

33. (Previously presented) The method of Claim 21, wherein the matrix is selected from the group consisting of poly(lactide-co-glycolide) (PLGA) and Poly vinylpyrrolidone (PVP).

34. (Currently amended) The method of Claim 21, further comprising:
applying a solution of a barrier material prior to ~~applying~~ introducing the first homogenous solution, the barrier material forming a barrier to the passage of the therapeutic agent in the first homogenous solution to one side of the body of the implantable medical device.

35. (Previously presented) The method of Claim 21, wherein the second homogenous solution contains no therapeutic agent.

36. (Previously presented) The method of Claim 21, wherein the second homogeneous solution comprises the at least one therapeutic agent, and wherein a concentration of the at least one therapeutic agent in the second homogeneous solution is different from a concentration of the at least one therapeutic agent in the first

homogeneous solution.

37. (Previously presented) The method of Claim 36, further comprising:
applying successive homogeneous solutions to the matrix; and
solidifying the successive homogeneous solutions, thereby forming additional portions of the matrix, wherein the concentration of the at least one therapeutic agent in the matrix is different in the successive portions of the matrix.

38. (Withdrawn) A method of forming an implantable medical device configured to release at least one therapeutic agent therefrom, wherein the therapeutic agent is disposed in a matrix affixed to a body of the implantable medical device, and wherein a concentration of the at least one therapeutic agent in the matrix varies as a continuous gradient relative to a surface of the implantable medical device body, the method comprising:

forming a homogeneous solution comprising a polymeric binder and a solvent;
evaporating the solvent in the homogeneous solution, thereby forming a matrix;
exposing the matrix to a solution comprising the therapeutic agent for a time sufficient to produce a partial diffusion of the therapeutic agent into the matrix such that the concentration of the therapeutic agent varies in the matrix; and
affixing the matrix to the implantable medical device body.

39. (Withdrawn) The method of Claim 38, wherein the matrix is affixed to the implantable medical device body by placing the matrix into the body prior to immersing the matrix in the solution comprising the therapeutic agent.

40. (Withdrawn) The method of Claim 38, wherein the matrix is affixed to the implantable medical device body by placing the matrix into a recess in the implantable

medical device body.

41. (Withdrawn) The method of Claim 38, wherein the matrix is affixed to the implantable medical device body by placing the matrix into an opening passing through the implantable medical device body.

42. (Withdrawn) The method of Claim 38, wherein the matrix is affixed to the implantable medical device body by disposing the homogeneous solution comprising a polymeric binder and a solvent into an opening and then evaporating the solvent.

43. (Withdrawn) The method of Claim 38, wherein the matrix is affixed to the implantable medical device body by coating a surface of the implantable medical device with the matrix.

44. (Withdrawn) A method for treating a patient by local delivery of at least one therapeutic agent, the method comprising:

delivering an implantable medical device into the body of a patient, the implantable medical device having a matrix affixed to a body of the implantable medical device with concentration of the at least one therapeutic agent in the matrix varying as a gradient relative to a surface of the body matrix with a high concentration of the therapeutic agent spaced a distance from the matrix surface and a low concentration of the therapeutic agent at the matrix surface; and

delivering the therapeutic agent at a release rate and over an administration period determined by the gradient of therapeutic agent in the matrix.

45-47. (Canceled).

48. (Previously presented) The method of Claim 21, wherein the implantable medical device is a stent.

49. (Withdrawn) The method of Claim 38, wherein the implantable medical device is a stent.

50. (Withdrawn) The method of Claim 44, wherein the implantable medical device is a stent.

51. (Withdrawn) The method of Claim 50, wherein the matrix is affixed to the stent by placing the matrix in a recess in the stent.